



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

But this is the same chance as that which was found above in different terms. Equating the two, we have

$$\frac{p! m(abc\dots)}{l^p} = \frac{n!(p-1)!}{(n-p)!(n+p-1)!}.$$

$$\text{Hence, } m(abc\dots) = \frac{l^p \cdot n!(p-1)!}{p!(n-p)!(n+p-1)!}.$$

For the case $p=n$ this value reduces to $\frac{l^n(n-1)!}{(2n-1)!}$, the mean value of the product of the n segments of the line. This result agrees with that found by Mr. Crofton in the solution of the latter problem in the *Encyclopedia Britannica*, Vol. XIX, p. 784.

This solution should have appeared instead of the solution published last month. Both solutions were received before the last issue went to press. By an oversight the defective solution got in with the material for publication. Ed. F.

170. Proposed by LON C. WALKER, Santa Barbara, Cal.

Find the area of a triangle formed by drawing a line at random through each of three points taken at random within a given triangle.

No solution has been received.

174. Proposed by J. EDWARD SANDERS, Reinersville, Ohio.

A chord of length C is drawn at random in a given ellipse. What is the average area of the segment cut off by the chord?

Solution by G. B. M. ZERR, A. M., Ph. D., Parsons, W. Va.

Refer the ellipse to conjugate diameters so that $x^2/m^2 + y^2/n^2 = 1$ is its equation.

$$\begin{aligned} \text{Area of segment} &= \frac{2n}{m} \sin \omega \int_{(m/2n)\sqrt{(4n^2-c^2)}}^m \sqrt{(m^2-x^2)} dx \\ &= mns \sin \omega \left[\cos^{-1} \left(\frac{\sqrt{(4n^2-c^2)}}{2n} \right) - \frac{c \sqrt{(4n^2-c^2)}}{4n^2} \right] \\ &= ab \left[\cos^{-1} \left(\frac{\sqrt{(4n^2-c^2)}}{2n} \right) - \frac{c \sqrt{(4n^2-c^2)}}{4n^2} \right] = A. \end{aligned}$$

ω is the inclination of m, n .

$$\text{Average area is } \Delta = \frac{\int_b^a A dn}{\int_b^a dn}.$$